

4

DTIC FILE COPY

AD-A196 902

OFFICE OF NAVAL RESEARCH

Contract N00014-86-K-0043

TECHNICAL REPORT No. 71

The coefficient of first viscosity via three-phonon
processes in bulk liquid helium

by

C. I. Um, C. W. Jun, W. H. Kahng and Thomas F. George

Prepared for Publication

in

Physical Review B

Departments of Chemistry and Physics
State University of New York at Buffalo
Buffalo, New York 14260

June 1988

Reproduction in whole or in part is permitted for any purpose of the
United States Government.

This document has been approved for public release and sale;
its distribution is unlimited.

DTIC
ELECTE
JUN 28 1988
S E D

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE

REPORT DOCUMENTATION PAGE				Form Approved OMB No. 0704-0188	
1a. REPORT SECURITY CLASSIFICATION Unclassified			1b. RESTRICTIVE MARKINGS		
2a. SECURITY CLASSIFICATION AUTHORITY			3. DISTRIBUTION/AVAILABILITY OF REPORT Approved for public release; distribution unlimited		
2b. DECLASSIFICATION/DOWNGRADING SCHEDULE					
4. PERFORMING ORGANIZATION REPORT NUMBER(S) UBUFFALO/DC/88/TR-71			5. MONITORING ORGANIZATION REPORT NUMBER(S)		
6a. NAME OF PERFORMING ORGANIZATION Depts. Chemistry & Physics State University of New York		6b. OFFICE SYMBOL (If applicable)	7a. NAME OF MONITORING ORGANIZATION		
6c. ADDRESS (City, State, and ZIP Code) Fronczak Hall, Amherst Campus Buffalo, New York 14260			7b. ADDRESS (City, State, and ZIP Code) Chemistry Program 800 N. Quincy Street Arlington, Virginia 22217		
8a. NAME OF FUNDING/SPONSORING ORGANIZATION Office of Naval Research		8b. OFFICE SYMBOL (If applicable)	9. PROCUREMENT INSTRUMENT IDENTIFICATION NUMBER Contract N00014-86-K-0043		
8c. ADDRESS (City, State, and ZIP Code) Chemistry Program 800 N. Quincy Street Arlington, Virginia 22217			10. SOURCE OF FUNDING NUMBERS		
			PROGRAM ELEMENT NO.	PROJECT NO.	TASK NO.
					WORK UNIT ACCESSION NO.
11. TITLE (Include Security Classification) The coefficient of first viscosity via three-phonon processes in bulk liquid helium					
12. PERSONAL AUTHOR(S) C. I. Um, C. W. Jun, W. H. Kahng and Thomas F. George					
13a. TYPE OF REPORT		13b. TIME COVERED FROM _____ TO _____		14. DATE OF REPORT (Year, Month, Day) June 1988	
				15. PAGE COUNT 15	
16. SUPPLEMENTARY NOTATION Prepared for publication in Physical Review B					
17. COSATI CODES			18. SUBJECT TERMS (Continue on reverse if necessary and identify by block number)		
FIELD	GROUP	SUB-GROUP	BULK LIQUID HELIUM, THEORETICAL STUDY		
			FIRST VISCOSITY, INVERSE TEMPERATURE DEPENDENCE		
			COEFFICIENT, THREE-PHONON PROCESSES.		
19. ABSTRACT (Continue on reverse if necessary and identify by block number) The contribution of three-phonon processes to the coefficient of first viscosity in bulk liquid helium is evaluated explicitly as a function of temperature, which is shown to have a T^{-1} dependence.					
20. DISTRIBUTION/AVAILABILITY OF ABSTRACT <input checked="" type="checkbox"/> UNCLASSIFIED/UNLIMITED <input checked="" type="checkbox"/> SAME AS RPT. <input type="checkbox"/> DTIC USERS			21. ABSTRACT SECURITY CLASSIFICATION Unclassified		
22a. NAME OF RESPONSIBLE INDIVIDUAL Dr. David L. Nelson			22b. TELEPHONE (Include Area Code) (202) 696-4410		22c. OFFICE SYMBOL

DD Form 1473, JUN 86

Previous editions are obsolete.

SECURITY CLASSIFICATION OF THIS PAGE

UNCLASSIFIED

**The coefficient of first viscosity via three-phonon
processes in bulk liquid helium**

C. I. Um, C. W. Jun and W. H. Kahng
Department of Physics, College of Science
Korea University
Seoul 136, Korea

Thomas F. George*
Departments of Chemistry and Physics
239 Fronczak Hall
State University of New York at Buffalo
Buffalo, New York 14260

The contribution of three-phonon processes to the coefficient of first viscosity in bulk liquid helium is evaluated explicitly as a function of temperature, which is shown to have a T^{-1} dependence.

PACS NO. 05.60 +W, 66.20.+d, 67.40 Pm.

Accession For	
NTIS GRA&I	<input checked="checked" type="checkbox"/>
DTIC TAB	<input type="checkbox"/>
Unannounced	<input type="checkbox"/>
Justification	
By _____	
Distribution/	
Availability Codes	
Dist	Avail and/or Special
A-1	

* To whom correspondence should be addressed.



1. Introduction

Since Landau and Khalatnikov's work¹⁾ on transport coefficients, i.e., the thermal conductivity and viscosities of liquid helium, Khalatnikov and Matveev²⁾ reexamined the transport coefficients due to contributions by phonon-roton scattering and noted that the results of Landau and Khalatnikov's work with a normal dispersion deviate from experimental data by as much as a factor of two. The correct anomalous dispersion indicates three-phonon processes (3PP) to be important near absolute zero temperature, and should thus be included in calculations of transport kinetic coefficients.

It is well known that the kinetic equation with temperature gradient for the thermal conductivity vanishes for a pure phonon gas, and thus the corresponding thermal conductivity becomes zero. However, the kinetic equation for the first viscosity does not vanish for a pure phonon gas near zero temperature, and 3PP become important. The purpose of this paper is to evaluate the contribution to the coefficient η_{ph} of first viscosity from 3PP which were not included in the work of Landau and Khalatnikov. For small momenta the three-dimensional dispersion relation has been evaluated explicitly by adopting a soft potential to give³⁾

$$\epsilon(p) = C_0 p [1 + \gamma p^2 - \delta p^3 + \dots] , \quad (1.1)$$

where C_0 is the velocity of sound at absolute zero temperature. For bulk liquid helium of density $2.18 \times 10^{-2} \text{ \AA}^{-3}$, we can use the following values⁴⁾ obtained to fit the experimental data⁵⁾ of the excitation energy spectrum:

$$\gamma \hbar^2 = 1.51 \text{ \AA}^2, \delta \hbar^3 = 3.25 \text{ \AA}^3, C_0 = 238.2 \text{ m/s}, u = 1.8 \quad ^{6)}$$

In Sec. 2 we consider phonon-phonon interactions in bulk liquid helium via second quantization to obtain the characteristic time for 3PP. We then solve the kinetic equation for the coefficient of first viscosity in Sec. 3. Finally, in Sec. 4 we discuss numerical results.

2. Phonon-phonon interaction in a three-dimensional quantum liquid

To consider the phonon-phonon interaction in a three-dimensional quantum liquid, we begin with the Hamiltonian H and the internal energy $E(\rho)$ ⁷⁾ for superfluid liquid ^4He , given by

$$H = \int d^3r \left[\frac{1}{2} \hat{v} \cdot \hat{p} \hat{v} + E(\rho) \right] , \quad (2.1)$$

$$E(\rho) = E(\rho_0) + \int d^3p \, n \epsilon(p) , \quad (2.2)$$

where \hat{v} and \hat{p} are the velocity and density operators expressed in terms of the annihilation operator \hat{a}_p and creation operator \hat{a}_p^\dagger , n is the number density of excitations, and \hat{p} is given by $\hat{p}_0 + \hat{p}'$, the constant and variable parts, respectively. Expanding the internal energy $E(\rho)$ as a function of ρ' and then expressing this expansion in terms of annihilation and creation operators, we obtain

$$H = H_0 + H_3 + H_4 + \dots \quad (2.3)$$

$$H_0 = \sum_p \left(\hat{a}_p^\dagger \hat{a}_p + \frac{1}{2} \right) \hbar \omega ,$$

$$\begin{aligned}
H_3 = & 4\pi^3 i \left(\frac{\hbar}{2V}\right)^{3/2} \sum_{p_1} \sum_{p_2} \sum_{p_3} \left[\left(\frac{C_0 p_3}{\rho_0 p_1 p_2}\right)^{1/2} (\vec{p}_1 \cdot \vec{p}_2) + \frac{\rho_0^2}{3C_0^2} \frac{\partial}{\partial \rho} \left(\frac{C_0}{\rho}\right)^2 \right. \\
& \times \left(\frac{C_0 p_1 p_2 p_3}{\rho_0}\right)^{1/2} \left. \left[\hat{a}_{p_1} \hat{a}_{p_2} \hat{a}_{p_3} \delta(\vec{p}_1 + \vec{p}_2 + \vec{p}_3) \exp(-i(\omega_1 + \omega_2 + \omega_3)t) \right. \right. \\
& \left. \left. + \hat{a}_{p_1} \hat{a}_{p_2}^\dagger \hat{a}_{p_3} \delta(\vec{p}_1 - \vec{p}_2 - \vec{p}_3) \exp(-i(\omega_1 - \omega_2 - \omega_3)t) + \dots \right] \right], \quad (2.4)
\end{aligned}$$

$$\begin{aligned}
H_4 = & \frac{(2\pi)^3}{4!} \left(\frac{\rho_0 \hbar}{2VC_0}\right)^2 \frac{\partial^2}{\partial \rho^2} \left(\frac{C_0^2}{\rho}\right) \sum_{p_1} \sum_{p_2} \sum_{p_3} \sum_{p_4} (p_1 p_2 p_3 p_4)^{1/2} \\
& \times \left[\hat{a}_{p_1} \hat{a}_{p_2} \hat{a}_{p_3} \hat{a}_{p_4} \delta(\vec{p}_1 + \vec{p}_2 + \vec{p}_3 + \vec{p}_4) \exp(-i(\omega_1 + \omega_2 + \omega_3 + \omega_4)t) \right. \\
& \left. + \hat{a}_{p_1} \hat{a}_{p_2}^\dagger \hat{a}_{p_3}^\dagger \hat{a}_{p_4} \delta(\vec{p}_1 - \vec{p}_2 - \vec{p}_3 + \vec{p}_4) \exp(-i(\omega_1 - \omega_2 - \omega_3 + \omega_4)t) + \dots \right], \quad (2.5)
\end{aligned}$$

where the other six terms to be included in Eq. (2.4) are given by combinations of $\hat{a}_{p_1}, \hat{a}_{p_2}, \hat{a}_{p_3}$ and $\hat{a}_{p_1}^\dagger, \hat{a}_{p_2}^\dagger, \hat{a}_{p_3}^\dagger$, and fourteen additional terms with the combinations of four creation and four annihilation operators appear in Eq. (2.5).

The nonzero matrix elements of \hat{a}_p and \hat{a}_p^\dagger are

$$\begin{aligned}
\langle n_p - 1 | \hat{a}_p | n_p \rangle &= -i(n_p)^{1/2} e^{i\omega t}, \\
\langle n_p | \hat{a}_p^\dagger | n_p - 1 \rangle &= i(n_p)^{1/2} e^{-i\omega t}.
\end{aligned} \quad (2.6)$$

The matrix element $\langle n_p | H_0 | n_p \rangle$ is $\sum_p (n_p + \frac{1}{2}) \hbar \omega$, where $\omega = c_0 \rho$ and H_0 is thus the Hamiltonian of a simple harmonic oscillator. H_3 and H_4 are the anharmonic perturbative Hamiltonians. Transitions between different states occur due to H_3 , H_4 and other higher-order perturbed Hamiltonians. The transition matrix element from state $|i\rangle$ to state $|f\rangle$ in second-order perturbation theory is given by

$$\langle f | H | i \rangle = \langle f | H_3 | i \rangle + \langle f | H_4 | i \rangle + \sum_j \frac{\langle f | H_3 | j \rangle \langle j | H_4 | i \rangle}{E_j - E_i} \quad (2.7)$$

According to Eqs. (2.6) and (2.4), the first term in Eq. (2.7) contributes only to 3PP, and the second and third terms are related to 4PP.

A finite lifetime of an elementary excitation in a quantum liquid originates from two processes. One is the collision between elementary excitations, and the other is the spontaneous decay of a phonon into two, three or more phonons. Since the collision probability tends to zero at very low temperatures, the collision process will become less important and negligible. At very low temperatures, below 0.6 K, the main excitations in liquid ^4He are phonons. Therefore, we may consider the decay of the phonon into two phonons only, i.e., 3PP,⁸⁾ consisting of two processes: the direct process of emission of a phonon, $\vec{p}_3, \vec{p}_1 = \vec{p}_2 + \vec{p}_3$, and the reverse process of absorption of a phonon, $\vec{p}_3, \vec{p}_2 + \vec{p}_3 = \vec{p}_1$. The differential decay probability per unit time in three dimensions is given by

$$d\omega = \frac{2\pi}{\hbar} |\langle f | H_3 | i \rangle|^2 \delta(\epsilon_f - \epsilon_i) \frac{v^2 d\vec{p}_2 d\vec{p}_3}{(2\pi\hbar)^6} \quad (2.8)$$

Since the matrix elements in both processes are given by

$$\begin{aligned} \langle f | H_3 | i \rangle_D = & \frac{3!}{2} \frac{(2\pi\hbar)^3}{(2V)^{3/2}} \delta(\vec{p}_1 - \vec{p}_2 - \vec{p}) \left(\frac{C_0}{\rho_0} \frac{p}{p_1 p_2} \right)^{1/2} (\vec{p}_1 \cdot \vec{p}_2) \\ & + \frac{\rho_0^2}{3C_0^2} \frac{\partial}{\partial \rho} \left(\frac{C_0^2}{\rho} \right) \left(\frac{C_0 p_1 p_2 p}{\rho_0} \right)^{1/2} \sqrt{n_{p_1} (n_{p_2} + 1) (n_p + 1)} , \end{aligned} \quad (2.9)$$

$$\begin{aligned} \langle f | H_3 | i \rangle_R = & \frac{3!}{2} \frac{(2\pi\hbar)^3}{(2V)^{3/2}} \delta(\vec{p}_1 - \vec{p}_2 - \vec{p}) \left(\frac{C_0}{\rho_0} \frac{p}{p_1 p_2} \right)^{1/2} (\vec{p}_1 \cdot \vec{p}_2) \\ & + \frac{\rho_0^2}{3C_0^2} \frac{\partial}{\partial \rho} \left(\frac{C_0^2}{\rho} \right) \left(\frac{C_0 p_1 p_2 p}{\rho_0} \right)^{1/2} \sqrt{(n_{p_1} + 1) n_{p_2} n_p} , \end{aligned} \quad (2.10)$$

where n_p is the distribution function of phonons with momentum p , the total decay probabilities per unit time of the two processes are

$$\omega_D = \frac{\pi C_0}{2\hbar \rho_0} (u+1)^2 \int \frac{d\vec{p}_2}{(2\pi\hbar)^3} p_1 p_2 p n_{p_1} (n_{p_2} + 1) (n_p + 1) \delta(\epsilon_f - \epsilon_i) , \quad (2.11)$$

$$\omega_R = \frac{\pi C_0}{2\hbar \rho_0} (u+1)^2 \int \frac{d\vec{p}_2}{(2\pi\hbar)^3} p_1 p_2 p (n_{p_1} + 1) n_{p_2} n_p \delta(\epsilon_f - \epsilon_i) , \quad (2.12)$$

where u is the Grüneisen constant given by $(\rho_0/C_0)(\partial C_0/\partial \rho_0)$.

The collision integral, which is the difference between the differential decay probability per unit time for the direct and reverse processes, can be written as

$$J_{3PP} = - \frac{\pi C_0}{2\hbar \rho_0} (u+1)^2 \int \frac{d\vec{p}_2}{(2\pi\hbar)^3} p_1 p_2 p \delta n(n_{p_1_0} - n_{p_2_0}) \delta(\epsilon_f - \epsilon_i) \quad (2.13)$$

where $\delta n = n_p - n_{p_0}$, with n_{p_0} as the distribution function of phonons with momentum p in the equilibrium state. Since the collision integral $J_{3PP}(n)$ is given by $-(n - n_0)/\tau_{3PP}$, we have

$$\tau_{3PP}^{-1} = \frac{\pi C_0}{2\hbar \rho_0} (u+1)^2 \int \frac{d\vec{p}_2}{(2\pi\hbar)^3} p_1 p_2 p (n_{p_1_0} - n_{p_2_0}) \delta(\epsilon_f - \epsilon_i) \quad (2.14)$$

The three-phonon processes are small-angle processes, and thus p_1 and p_2 are much larger than p_3 . Hence we may use the following relations to evaluate Eq. (2.14):

$$n_{p_1_0} - n_{p_2_0} = \left(\frac{\partial n_{p_1_0}}{\partial \omega_1} \right) \omega \quad (2.15)$$

$$\omega_2 - \omega_1 \approx - \left(\frac{\partial \omega_1}{\partial p_1} \right) p_3 = -\omega \cos \theta \quad (2.16)$$

Substitution of Eqs. (2.15)-(2.16) into Eq. (2.14) and integration of Eq. (2.14) over momentum space yields

$$\frac{1}{\tau_{3PP}} = \omega \frac{\pi^3 (u+1)^2 k_B^4 T^4}{30\hbar^3 \rho_0 C_0^5} \left[1 - \frac{60\pi^2}{7} \gamma \left(\frac{k_B T}{C_0} \right)^2 \right] \quad (2.17)$$

3. Viscosity

In this section we shall evaluate the coefficient of first viscosity. We first consider the macroscopic motion of liquid helium, which does not depend on time, with velocity \vec{v} in the z-direction and the gradient directed along x-axis. Then the kinetic equation becomes

$$\vec{v} \cdot \vec{\nabla} n = v_x \frac{\partial n}{\partial x} = J(n) \quad (3.1)$$

The phonon distribution function in the motion of liquid helium with velocity \vec{v} is given by

$$n_o = [\exp(\frac{Cp - \vec{v} \cdot \vec{p}}{k_B T}) - 1]^{-1} \quad (3.2)$$

where C is the phonon velocity (equal to C_o at zero temperature). Substitution of Eq. (3.2) into (3.1) yields

$$n_o(n_o + 1) \frac{Cp}{k_B T} \frac{\partial v}{\partial x} \cos\theta \sin\theta \cos\phi = J(n) \quad (3.3)$$

where here θ is the angle between \vec{p} and the z-axis. Since 4PP establish energy in the equilibrium state and the distribution function of phonons is related to the variation of phonons, which characterize the chemical potential α , the distribution function becomes

$$n = [\exp(\alpha' + \frac{pC}{k_B T} - 1)]^{-1} \quad (3.4)$$

Expanding Eq. (3.4) as a function of α' to obtain the deviation of n from n_0 and considering the linearity of the left-hand side in Eq. (3.3), we can write

$$\delta n = n - n_0 = -n_0(n_0+1)\alpha' \quad (3.5)$$

$$\alpha' = \alpha \cos\theta \sin\theta \cos\phi \quad (3.6)$$

Replacing $J(n)$ by $J_{3PP}(n)$ in Eq. (3.3) and integrating over momentum space, we obtain

$$\int dp p^2 J_{3PP}(n) = \cos\theta \sin\theta \cos\phi \left(\frac{k_B T}{C_0}\right)^3 \frac{3!}{3!} \zeta(3) \frac{\partial v}{\partial x} \quad (3.7)$$

Since the collision integral $J_{3PP}(n)$ can be replaced by $-(n-n_0)/\tau_{3PP}$, we can use Eqs. (3.5)-(3.6) together with Eq. (2.17) and $\omega = pC_0/\hbar$ for the evaluation of collision integral:

$$\int dp p^2 J_{3PP}(n) = \frac{\pi^3 (u+1)^2 (k_B T)^4}{30 \hbar^4 \rho_0 C_0^4} \left(\frac{k_B T}{C_0}\right)^4 \frac{3!}{3!} \zeta(3) \alpha \cos\theta \sin\theta \cos\phi \quad (3.8)$$

From comparison of Eqs. (3.7) and (3.8) we have

$$\alpha = \frac{30 \hbar^4 \rho_0 C_0^5}{\pi^3 (u+1)^2 (k_B T)^5} \frac{\partial v}{\partial x} = A \frac{\partial v}{\partial x} \quad (3.9)$$

The coefficient of first viscosity is obtained from the evaluation of the stress tensor associated with 3PP given by

$$\begin{aligned}
\sigma_{xz} &= -\eta_{ph} \frac{\partial v}{\partial x} = \int \frac{d\vec{p}}{(2\pi\hbar)^3} p_z v_x (n - n_0) \\
&= \int dp \frac{p^2 \sin\theta d\theta d\phi}{(2\pi\hbar)^3} p \cos\theta C_0 \sin\theta \cos\phi [-n_0(n_0+1) A \frac{\partial v}{\partial x} \cos\theta \sin\theta \cos\phi]
\end{aligned} \quad (3.10)$$

Performing the integral, Eq. (3.10) becomes

$$\sigma_{xz} = -\frac{3!5(3)}{30\pi^2\hbar^3} \frac{(k_B T)^4}{C_0^3} A \frac{\partial v}{\partial x} \quad (3.11)$$

We can get the coefficient of first viscosity by comparing Eq. (3.11) with the first expression in Eq. (3.10):

$$\eta_{ph}(T) = \frac{3!5(3)\hbar\rho_0 C_0^2}{\pi^5 k_B (u+1)^2} \frac{1}{T} = 1.904 \times 10^{-7} \frac{1}{T} \quad (3.12)$$

We note that the coefficient of first viscosity^{1),9)} at temperatures below 0.9 K is given by

$$\eta_{ph}(T) = 3.50 \times 10^{-9} T^{-1/2} e^{\Delta/k_B T} (1 + 2.15 \times 10^{-5} T^{9/2} e^{\Delta/k_B T})^{-1}, \quad T < 0.9 \text{ K} \quad (3.13)$$

4. Numerical Results and discussion

In previous sections we have evaluated the lifetime and the contribution of first viscosity. At relatively low temperatures (< 0.9 K), Eq. (3.13) reduces approximately to

$$\eta_{ph} \sim 1.5 \times 10^{-4} \frac{1}{T^5} .$$

However, as the temperature approaches absolute zero, the roton density becomes zero, and thus the contribution from the roton-phonon scattering to viscosity will vanish so that the main contribution comes from the 3PP. Therefore, the temperature dependence is changed from T^{-5} to T^{-1} . Figure 1 illustrates the coefficient $\eta_{ph}(T)$ given by Eq. (3.12). The coefficient increases slowly, and near zero temperature it increases very rapidly and finally diverges. In conclusion, we point out that in consideration of the 3PP originating from the anomalous excitation spectrum at low momenta, the temperature variation of first viscosity depends on T^{-1} .

Acknowledgments

The authors are very thankful to Professor J. S. Kang for a critical reading of the manuscript. This research was supported in part by a grant to Korea University from KOSEF under Contract No. 862-0203-011-2, and in part by the Office of Naval Research and Air Force Office of Scientific Research (AFSC), under Contract F49620-86-C-0009, with the Research Foundation of the State University of New York. The United States Government is authorized to reproduce and distribute reprints for governmental purposes notwithstanding any copyright notation hereon.


References

1. L. D. Landau and I. M. Khalatnikov, Zh. Eksp. Teor. Fiz. 19, 637, 709 (1949).
2. I. M. Khalatnikov and Yu. A. Matveev, J. Low Temp. Phys. 51, 99 (1983).
3. C. I. Um, W. H. Kahng and A. Isihara, Physica 100B, 74 (1980); 114B, 191 (1982).
4. C. I. Um, S. T. Choh and A. Isihara, Phys. Rev. B 23, 4498 (1981).
5. D. Henshaw and A. Woods, Phys. Rev. 121, 266 (1961).
6. K. K. Singh and J. Prakash, Phys. Rev. B 17, 1253 (1978).
7. R. N. Bhatt and W. L. McMillan, Phys. Rev. A 10, 1591 (1974).
8. R. C. Dynes and V. Narayanamurti, Phys. Rev. B 12, 1720 (1975); H. J. Maris, Rev. Mod. Phys. 49, 341 (1977).
9. J. Wilks, The Properties of Liquid and Solid Helium (Oxford University Press, Oxford, 1967), Chapt. 7, p. 177; K. H. Bennemann and J. B. Ketterson, The Physics of Liquid and Solid Helium (Wiley, New York, 1976), Part 1, Chapt. 1, p. 71.

Figure Caption

Figure 1. Viscosity coefficient due to the contribution from 3PP.

Fig. 1



TECHNICAL REPORT DISTRIBUTION LIST, GEN

	<u>No. Copies</u>		<u>No. Copies</u>
Office of Naval Research Attn: Code 1113 800 N. Quincy Street Arlington, Virginia 22217-5000	2	Dr. David Young Code 334 NORDA NSTL, Mississippi 39529	1
Dr. Bernard Douda Naval Weapons Support Center Code 50C Crane, Indiana 47522-5050	1	Naval Weapons Center Attn: Dr. Ron Atkins Chemistry Division China Lake, California 93555	1
Naval Civil Engineering Laboratory Attn: Dr. R. W. Drisko, Code L52 Port Hueneme, California 93401	1	Scientific Advisor Commandant of the Marine Corps Code RD-1 Washington, D.C. 20380	1
Defense Technical Information Center Building 5, Cameron Station Alexandria, Virginia 22314	12 high quality	U.S. Army Research Office Attn: CRD-AA-IP P.O. Box 12211 Research Triangle Park, NC 27709	1
DTNSRDC Attn: Dr. H. Singerman Applied Chemistry Division Annapolis, Maryland 21401	1	Mr. John Boyle Materials Branch Naval Ship Engineering Center Philadelphia, Pennsylvania 19112	1
Dr. William Tolles Superintendent Chemistry Division, Code 6100 Naval Research Laboratory Washington, D.C. 20375-5000	1	Naval Ocean Systems Center Attn: Dr. S. Yamamoto Marine Sciences Division San Diego, California 91232	1
		Dr. David L. Nelson Chemistry Division Office of Naval Research 800 North Quincy Street Arlington, Virginia 22217	1

ABSTRACTS DISTRIBUTION LIST, 056/625/629

Dr. J. E. Jensen
Hughes Research Laboratory
3011 Malibu Canyon Road
Malibu, California 90265

Dr. C. B. Harris
Department of Chemistry
University of California
Berkeley, California 94720

Dr. J. H. Weaver
Department of Chemical Engineering
and Materials Science
University of Minnesota
Minneapolis, Minnesota 55455

Dr. F. Kutzler
Department of Chemistry
Box 5055
Tennessee Technological University
Cookeville, Tennessee 38501

Dr. A. Reisman
Microelectronics Center of North Carolina
Research Triangle Park, North Carolina
27709

Dr. D. DiLella
Chemistry Department
George Washington University
Washington D.C. 20052

Dr. M. Grunze
Laboratory for Surface Science and
Technology
University of Maine
Orono, Maine 04469

Dr. R. Reeves
Chemistry Department
Rensselaer Polytechnic Institute
Troy, New York 12181

Dr. J. Butler
Naval Research Laboratory
Code 6115
Washington D.C. 20375-5000

Dr. Steven M. George
Stanford University
Department of Chemistry
Stanford, CA 94305

Dr. L. Interante
Chemistry Department
Rensselaer Polytechnic Institute
Troy, New York 12181

Dr. Mark Johnson
Yale University
Department of Chemistry
New Haven, CT 06511-8118

Dr. Irvin Heard
Chemistry and Physics Department
Lincoln University
Lincoln University, Pennsylvania 19352

Dr. W. Knauer
Hughes Research Laboratory
3011 Malibu Canyon Road
Malibu, California 90265

Dr. K.J. Klaubunde
Department of Chemistry
Kansas State University
Manhattan, Kansas 66506

ABSTRACTS DISTRIBUTION LIST, 056/625/629

Dr. G. A. Somorjai
Department of Chemistry
University of California
Berkeley, California 94720

Dr. J. Murday
Naval Research Laboratory
Code 6170
Washington, D.C. 20375-5000

Dr. J. B. Hudson
Materials Division
Rensselaer Polytechnic Institute
Troy, New York 12181

Dr. Theodore E. Madey
Surface Chemistry Section
Department of Commerce
National Bureau of Standards
Washington, D.C. 20234

Dr. J. E. Demuth
IBM Corporation
Thomas J. Watson Research Center
P.O. Box 218
Yorktown Heights, New York 10598

Dr. M. G. Lagally
Department of Metallurgical
and Mining Engineering
University of Wisconsin
Madison, Wisconsin 53706

Dr. R. P. Van Duyne
Chemistry Department
Northwestern University
Evanston, Illinois 60637

Dr. J. M. White
Department of Chemistry
University of Texas
Austin, Texas 78712

Dr. D. E. Harrison
Department of Physics
Naval Postgraduate School
Monterey, California 93940

Dr. R. L. Park
Director, Center of Materials
Research
University of Maryland
College Park, Maryland 20742

Dr. W. T. Peria
Electrical Engineering Department
University of Minnesota
Minneapolis, Minnesota 55455

Dr. Keith H. Johnson
Department of Metallurgy and
Materials Science
Massachusetts Institute of Technology
Cambridge, Massachusetts 02139

Dr. S. Sibener
Department of Chemistry
James Franck Institute
5640 Ellis Avenue
Chicago, Illinois 60637

Dr. Arnold Green
Quantum Surface Dynamics Branch
Code 3817
Naval Weapons Center
China Lake, California 93555

Dr. A. Wold
Department of Chemistry
Brown University
Providence, Rhode Island 02912

Dr. S. L. Bernasek
Department of Chemistry
Princeton University
Princeton, New Jersey 08544

Dr. W. Kohn
Department of Physics
University of California, San Diego
La Jolla, California 92037

ABSTRACTS DISTRIBUTION LIST, 056/625/629

Dr. F. Carter
Code 6170
Naval Research Laboratory
Washington, D.C. 20375-5000

Dr. Richard Colton
Code 6170
Naval Research Laboratory
Washington, D.C. 20375-5000

Dr. Dan Pierce
National Bureau of Standards
Optical Physics Division
Washington, D.C. 20234

Dr. R. Stanley Williams
Department of Chemistry
University of California
Los Angeles, California 90024

Dr. R. P. Messmer
Materials Characterization Lab.
General Electric Company
Schenectady, New York 22217

Dr. Robert Gomer
Department of Chemistry
James Franck Institute
5640 Ellis Avenue
Chicago, Illinois 60637

Dr. Ronald Lee
R301
Naval Surface Weapons Center
White Oak
Silver Spring, Maryland 20910

Dr. Paul Schoen
Code 6190
Naval Research Laboratory
Washington, D.C. 20375-5000

Dr. John T. Yates
Department of Chemistry
University of Pittsburgh
Pittsburgh, Pennsylvania 15260

Dr. Richard Greene
Code 5230
Naval Research Laboratory
Washington, D.C. 20375-5000

Dr. L. Kesmodel
Department of Physics
Indiana University
Bloomington, Indiana 47403

Dr. K. C. Janda
University of Pittsburgh
Chemistry Building
Pittsburg, PA 15260

Dr. E. A. Irene
Department of Chemistry
University of North Carolina
Chapel Hill, North Carolina 27514

Dr. Adam Heller
Bell Laboratories
Murray Hill, New Jersey 07974

Dr. Martin Fleischmann
Department of Chemistry
University of Southampton
Southampton SO9 5NH
UNITED KINGDOM

Dr. H. Tachikawa
Chemistry Department
Jackson State University
Jackson, Mississippi 39217

Dr. John W. Wilkins
Cornell University
Laboratory of Atomic and
Solid State Physics
Ithaca, New York 14853

ABSTRACTS DISTRIBUTION LIST, 056/625/629

Dr. R. G. Wallis
Department of Physics
University of California
Irvine, California 92664

Dr. D. Ramaker
Chemistry Department
George Washington University
Washington, D.C. 20052

Dr. J. C. Hemminger
Chemistry Department
University of California
Irvine, California 92717

Dr. T. F. George
Chemistry Department
University of Rochester
Rochester, New York 14627

Dr. G. Rubloff
IBM
Thomas J. Watson Research Center
P.O. Box 218
Yorktown Heights, New York 10598

Dr. Horia Metiu
Chemistry Department
University of California
Santa Barbara, California 93106

Dr. W. Goddard
Department of Chemistry and Chemical
Engineering
California Institute of Technology
Pasadena, California 91125

Dr. P. Hansma
Department of Physics
University of California
Santa Barbara, California 93106

Dr. J. Baldeschwieler
Department of Chemistry and
Chemical Engineering
California Institute of Technology
Pasadena, California 91125

Dr. J. T. Keiser
Department of Chemistry
University of Richmond
Richmond, Virginia 23173

Dr. R. W. Plummer
Department of Physics
University of Pennsylvania
Philadelphia, Pennsylvania 19104

Dr. E. Yeager
Department of Chemistry
Case Western Reserve University
Cleveland, Ohio 44106

Dr. N. Winograd
Department of Chemistry
Pennsylvania State University
University Park, Pennsylvania 16802

Dr. Roald Hoffmann
Department of Chemistry
Cornell University
Ithaca, New York 14853

Dr. A. Steckl
Department of Electrical and
Systems Engineering
Rensselaer Polytechnic Institute
Troy, New York 12181

Dr. G.H. Morrison
Department of Chemistry
Cornell University
Ithaca, New York 14853